



Purina Mills



Powerful Performance. Winning Results.



Showpig Grower L200

Feed Honor® Show Chow® Showpig Grower L200 to growing barrows and gilts as the sole ration.

Honor® Show Chow® Showpig Grower L200 feed is supported by Purina Mills® Animal Nutrition. Get your best animals on Honor® Show Chow® Showpig Grower L200 today so they can reach their full potential.

FEATURES	BENEFITS
Formulated for optimum growth and maximum muscle development	Helps unlock the pig's genetic potential
High energy formula for today's showpigs	For optimum growth and maximum muscle deposition
Contains 19.5% crude protein	Helps to fuel muscle synthesis without the hard, ripped look
Medicated with Lincomycin	Helps support normal immune systems
High quality animal protein ingredients	Helps improve appearance of muscle, skin and hair

Visit our website to download this product form.



Regional Availability

Check your current price list for availability in your region, or contact your local Land O' Lakes Purina Feeds Dealer or Representative for more information.

For more information on this product, visit our website at www.BetterAnimalsShowSeries.com or call Land O' Lakes Purina Feeds Customer Service at 1-800-227-8941

www.BetterAnimalsShowSeries.com

Showpig Grower L200

GUARANTEED ANALYSIS

Crude Protein, %	(min)	19.5
Lysine, %	(min)	1.20
Crude Fat, %	(min)	5.00
Crude Fiber, %	(max)	3.00
Calcium, % (Ca)	(min)	0.60
Calcium, % (Ca)	(max)	1.10
Phosphorus, % (P)	(min)	0.70
Salt, % (Na Cl)	(min)	0.20
Salt, % (Na Cl)	(max)	0.70
Selenium, ppm (Se)	(min)	0.30
Zinc, ppm (Zn)	(min)	250.00

Medicated for reduction in the severity of swine Mycoplasmal pneumonia caused by *Mycoplasma hyopneumoniae*.

Active Drug Ingredient: Lincomycin (*as lincomycin hydrochloride*) 200 g/ton

FEEDING DIRECTIONS

Feed to growing swine as the sole ration for 21 days.

Caution:

Occasionally, swine fed lincomycin may within the first two days after the onset of treatment develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions have been self-correcting within five to eight days without discontinuing the lincomycin treatment. Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects. Not to be fed to swine that weigh more than 250 lbs.

Reminder:

Feed is perishable and should be stored in a clean, dry, well-ventilated area so it will remain fresh and palatable. DO NOT feed moldy or insect infested feed to animals as it may cause illness, abortion or death.

Fresh, clean water should be available at all times.